

it's **GO** time

VOXZOGO® (vosoritide) is a prescription medicine used to increase linear growth in children with achondroplasia and open growth plates.

Annika, 14 years old,
receiving VOXZOGO
as part of a clinical trial

VOXZOGO®
(vosoritide) for injection

IMPORTANT SAFETY INFORMATION

VOXZOGO may cause serious side effects including a temporary decrease in blood pressure in some patients. To reduce the risk of a decrease in blood pressure and associated symptoms (dizziness, feeling tired, or nausea), patients should eat a meal and drink 8 to 10 ounces of fluid within 1 hour before receiving VOXZOGO.

Please see Important Safety Information on page 4, as well as the included full Prescribing Information and Patient Information.

Important information for traveling with VOXZOGO

Keeping up with daily injections is important for getting the most from VOXZOGO® (vosoritide). While taking a trip may disrupt your routine, a little advance planning can ease the travel experience for you and your child.



Pack enough supplies for your daily injection to cover each day of the trip

Consider bringing additional medicine and supplies in case your time away is extended.

Use this checklist to pack treatment-related supplies

- ☐ ____ days of VOXZOGO in its original packaging
- ☐ ____ days of supplies (eg, alcohol swabs, bandages, rewards)
- ☐ Insulated travel container (with cold packs, if not already brought up to room temperature)
- ☐ Sharps disposal container (consider purchasing a travel size)
- ☐ Doctor's note explaining VOXZOGO and what it treats (for air travel; if traveling internationally, consider having it translated to ease the screening process)



Administer VOXZOGO around the same time each day

Consider setting a reminder to help you keep the same interval between doses. Be sure to adjust for traveling across time zones.



Travel may expose your medication to extreme temperatures, which could compromise its effectiveness

- ▶ Temperature excursions are permitted at 59°F to 86°F for 90 days
- ▶ Once VOXZOGO has been brought up to room temperature, do not return it to the refrigerator
- ▶ Consider using cold packs in an insulated travel container if you prefer to keep the product refrigerated

VOXZOGO[®]
(vosoritide) for injection

Be sure to review the VOXZOGO
Instructions for Use (IFU).





If you're driving to your destination:

- ▶ Make sure your insulated travel container with VOXZOGO stays out of direct sunlight during your car trip
- ▶ Do not leave it in an unattended car for an extended period of time



If you're flying domestically or internationally:

- ▶ Pack VOXZOGO in your carry-on luggage—do not pack it in your checked luggage!
- ▶ Inform a TSA agent that you have medically necessary liquids and equipment before going through security
- ▶ The effects of x-ray screening on VOXZOGO have not been studied. Let the TSA officer know you'd prefer a hand inspection for your medication and supplies

Check the [TSA website](#) for the most up-to-date information about traveling with medications, including any necessary documentation.



If your child is traveling alone:

- ▶ Use the VOXZOGO checklist in this guide and review the Instructions for Use and Patient Information to ensure your child has all the supplies needed
- ▶ Confirm that the person who will be administering VOXZOGO is well trained and comfortable with the injection process—if not, reach out to your BioMarin Clinical Coordinator for additional training



If you're traveling for longer than 30 days:

- ▶ Contact your specialty pharmacy for help with medication supply needed for extended travel and any temporary address changes

Reach out to your BioMarin Clinical Coordinator or Specialty Pharmacy for more information about traveling with VOXZOGO.

IMPORTANT SAFETY INFORMATION

What is VOXZOGO used for?

- VOXZOGO is a prescription medicine used to increase linear growth in children with achondroplasia and open growth plates (epiphyses).
- VOXZOGO is approved under accelerated approval based on an improvement in annualized growth velocity. Continued approval may be contingent upon verification and description of clinical benefit in confirmatory trials.

What is the most important safety information about VOXZOGO?

- VOXZOGO may cause serious side effects including a temporary decrease in blood pressure in some patients. To reduce the risk of a decrease in blood pressure and associated symptoms (dizziness, feeling tired, or nausea), patients should eat a meal and drink 8 to 10 ounces of fluid within 1 hour before receiving VOXZOGO.

What are the most common side effects of VOXZOGO?

- The most common side effects of VOXZOGO include injection site reactions (including redness, itching, swelling, bruising, rash, hives, and injection site pain), high levels of blood alkaline phosphatase shown in blood tests, vomiting, joint pain, decreased blood pressure, and stomach ache. These are not all the possible side effects of VOXZOGO. Ask your healthcare provider for medical advice about side effects, and about any side effects that bother the patient or that do not go away.

How is VOXZOGO taken?

- VOXZOGO is taken daily as an injection given under the skin, administered by a caregiver after a healthcare provider determines the caregiver is able to administer VOXZOGO. Do not try to inject VOXZOGO until you have been shown the right way by your healthcare provider. VOXZOGO is supplied with Instructions for Use that describe the steps for preparing, injecting, and disposing VOXZOGO. Caregivers should review the Instructions for Use for guidance and any time they receive a refill of VOXZOGO in case any changes have been made.
- Inject VOXZOGO 1 time every day, at about the same time each day. If a dose of VOXZOGO is missed, it can be given within 12 hours from the missed dose. After 12 hours, skip the missed dose and administer the next daily dose as usual.
- The dose of VOXZOGO is based on body weight. Your healthcare provider will adjust the dose based on changes in weight following regular check-ups.
- Your healthcare provider will monitor the patient's growth and tell you when to stop taking VOXZOGO if they determine the patient is no longer able to grow. Stop administering VOXZOGO if instructed by your healthcare provider.

What should you tell the doctor before or during taking VOXZOGO?

- Tell your doctor about all of the patient's medical conditions including
 - If the patient has heart disease (cardiac or vascular disease), or if the patient is on blood pressure medicine (anti-hypertensive medicine).
 - If the patient has kidney problems or renal impairment.
 - If the patient is pregnant or plans to become pregnant. It is not known if VOXZOGO will harm the unborn baby.
 - If the patient is breastfeeding or plans to breastfeed. It is not known if VOXZOGO passes into breast milk.
- Tell your doctor about all of the medicines the patient takes, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

You may report side effects to BioMarin at **1-866-906-6100**. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call **1-800-FDA-1088**.

Please see additional safety information in the full [Prescribing Information](#) and [Patient Information](#).